Asymptotics of Few-body Equations of Collision Theory

Vagner Jikia, Jemal Mebonia
I. Javakhishvili Tbilisi State University, Tbilisi, Georgia,
Email: v_ jikia@yahoo.com

Abstract. It is well known that in the case of three or more particles the Lipmann-Schwinger (LS) integral equation is not of Fredholm type. This is the consequence of the existence of bounded subsystems. By using identical transformations of the kernel of LS equation for certain classes of potentials Faddeev obtained Fredholm type integral equations for the three-body problem [1]. The motion of for bodies is described by equations of Yakubovsky and Alt-Grassberger-Sandhas-Khelashvili [2.3], which are obtained as a result of two subsequent transpormations of the kernel of LS equation. It has been shown that for $N > 4$, on the z complex plane wherever $\mathrm{Im}z \neq 0$, after $N-2$ iterations the kernel of N-particle integral equations of the collision theory became compact. For $\mathrm{Im}z = 0$ the compactness has not been proven yet [4]. In turned out that, in the case of $N \geq 3$, the mentioned equations have correct asymptotic form the derivation and detailed analysis of which is presented this paper.

Keywords and phrases: Collision theory, asymptotic integral equations, Unitary asymptotic solution.

arXiv:1403.4483v1 [math-ph] 18 Mar 2014

Few-body equations, general formalism

Let us consider a system of N particles in non-relativistic approximation. Interaction operator V in the approximation of two-particle forces is given as:

$$V = \sum_{\alpha} v_{\alpha}, \quad \alpha \equiv (mn); \quad m, n = 1, 2, \ldots, N; \ m < n, N \geq 3. \tag{1}$$

Here v_α are potentials of pair-interactions.

N-particle scattering T-matrix is determined by the LS equation:

$$T(z) = V + VG_0(z)\,T(z). \tag{2}$$

Here G_0 is the Green's function of N free particles:

$$G_0(z) = (z - H_0)^{-1}, \quad z = E_0 + i\varepsilon, \tag{3}$$

where H_0 is the operator of kinetic energy and E_0 is the energy of the system's free motion. G_0 can be presented as a sum of anti-Hermitian and Hermitian parts:

$$G_0(z) = G_1(z) + G_2(z), \tag{4}$$

$$G_1(z) = \tfrac{1}{2}\left(G_0(z) - G_0(\tilde{z})\right), \quad G_2(z) = \tfrac{1}{2}\left(G_0(z) + G_0(\tilde{z})\right). \tag{5}$$

Here $\tilde{z}$ stands for the complex conjugate of the z parameter.

To solve equation (2) formally one applies the standard Faddeev approach and obtains:

$$T(z) = \sum_{\alpha} T^{\alpha}(z), \tag{6}$$

where auxillary operators T^α are define by the following system of equations:

$$T^{\alpha}(z) = T_{\alpha}(z) + T_{\alpha}(z)\,G_0(z) \sum_{\beta \neq \alpha} T^{\beta}(z), \tag{7}$$

and for two-particle operators T_α one can write:

$$T_{\alpha}(z) = v_{\alpha} + v_{\alpha} G_0(z)\,T_{\alpha}(z). \tag{8}$$

Equation (2) can be also solved formally by applying the Heitler formalism of separation [6] resulting in:

$$T(z) = K(z) + K(z)\, G_1(z)\, T(z), \tag{9}$$

$$K(z) = V + V G_2(z)\, K(z). \tag{10}$$

From the Heitler equation (9) T can be expressed in terms of a Hermitian operator K which in turn is determined by Eq. (10). If Eqs. (9) and (10) are correct equations (the two-particle case) then the T operator constructed this way satisfies the corresponding condition of unitarity [6] for any approximation of K.

Similarly to Eq. (2) let us look for the solution of Eq. (10) in the form of a sum:

$$K(z) = \sum_{\alpha} K^{\alpha}(z), \tag{11}$$

where auxiliary operators K^{α} are given by equation:

$$K^{\alpha}(z) = K_{\alpha}(z) + K_{\alpha}(z)\, G_2(z) \sum_{\beta \neq \alpha} K^{\beta}(z), \tag{12}$$

and for two particle operators K_{α} we have:

$$K_{\alpha}(z) = \upsilon_{\alpha} + \upsilon_{\alpha} G_2(z)\, K_{\alpha}(z). \tag{13}$$

It is well known that Eqs. (7) and (12) are not compact in general.

Below we show that by using the Heitler formalism it is possible to find the correct asymptotic (high energy) form of the system of multi-particle integral equations (7).

Multi-particle equations in asymptotics

Using the behaviour of Eq. (12) at sufficiently high energies: $\mathrm{Re}(z) \gg E^B_{\min}$ ($E^B_{\min}$ denotes the absolute magnitude of the minimum of all allowed binding energies in the system to be considered), let us restrict Eq. (9) to its high-energy approximation and using the obtained result let us rewrite Eq. (7) in asymptotics.

It is well-known that the square of the of Eq. (12) can be expressed as a sum of terms of the following form:

$$K_{\alpha}(z)\, G_2(z)\, K_{\beta}(z)\, G_2(z), \quad \alpha \neq \beta, \tag{14}$$

which in the approximation of Eq. (36) can be negleqted. This is why the kernel of Eq. (12) at sufficiently high energies is square integrable so that the corresponding Neuman series converges. To obtain the desired system of equations let us keep the first order asymptotic terms in Eq. (12) (K-matrix impulse approximation, see [7]):

$$K^{\alpha}(z) = K_{\alpha}(z), \quad K(z) = \sum_{\alpha} K_{\alpha}(z). \tag{15}$$

We substitute the K operator of Eqs. (15) into Eq. (9) and obtain:

$$T(z) = \sum_{\alpha} K_{\alpha}(z) + \sum_{\alpha} K_{\alpha}(z) G_1(z)\, T(z). \tag{16}$$

Equation (16) represents the Heitler equation in the K-matrix impulse approximation. It is defined in the kinematic area: $\mathrm{Re}(z) \gg E^B_{\min}$. Let us demonstrate that by using the standard Faddeev approach Eq. (16) reduces to the desired system of integral equations. For this purpose we decompose T in the form of Eq. (6) where, according to Eq. (16)

$$T^{\alpha}(z) = K_{\alpha}(z) + K_{\alpha}(z)\, G_1(z)\, T(z) \tag{17}$$

and insert Eq. (6) into Eq. (17). Inverting the diagonal elements in the resulting expression we obtain:

$$\left(1-K_{\alpha}\left(z\right)G_{1}\left(z\right)\right)^{-1}K_{\alpha}\left(z\right)+\left(1-K_{\alpha}\left(z\right)G_{1}\left(z\right)\right)^{-1}K_{\alpha}\left(z\right)G_{1}\left(z\right)\sum_{\beta\neq\alpha}T^{\beta}\left(z\right). \tag{18}$$

Using two-particle equations:

$$T_{\alpha}\left(z\right)=K_{\alpha}\left(z\right)+K_{\alpha}\left(z\right)G_{1}\left(z\right)T_{\alpha}\left(z\right), \tag{19}$$

one can write:

$$K_{\alpha}\left(z\right)=\left(1+T_{\alpha}\left(z\right)G_{1}\left(z\right)\right)^{-1}T_{\alpha}\left(z\right). \tag{20}$$

Inserting Eq. (20) into Eq. (18) we obtain:

$$T^{\alpha}\left(z\right)=T_{\alpha}\left(z\right)+T_{\alpha}\left(z\right)G_{1}\left(z\right)\sum_{\beta\neq\alpha}T^{\beta}\left(z\right). \tag{21}$$

Eq. (21) represents asymptotic $\left(\mathrm{Re}z\gg E_{\min}^{B}\right)$ form of the system (7) of N-particle integral equations, which indicates that at high energies in the Green's function G_0 entering in Eq. (7) its Hermitian part can be negleqted (for the two-particle case see Ref. [8]). Equation (21) is a system of equations for auxiliary operators T^{α} in the K-matrix impulse approximation. According to the estimation (see appendix, Eq. (38))

$$\left\|T_{\alpha}\left(z\right)G_{1}\left(z\right)T_{\beta}\left(z\right)G_{1}\left(z\right)\right\|=O\left(\varepsilon^{2}\right)\approx0,\quad\alpha\neq\beta, \tag{22}$$

the kernel of Eq. (21) is square integrable and therefore it is the fredholm type system of integral equations, iterative series of which with the accuracy of Eq. (22) converges to the following finite sum:

$$T^{\alpha}\left(z\right)=T_{\alpha}\left(z\right)+T_{\alpha}\left(z\right)G_{1}\left(z\right)\sum_{\beta\neq\alpha}T_{\beta}\left(z\right),\quad\mathrm{Re}\left(z\right)\gg E_{\min}^{B}. \tag{23}$$

It can be shown that the specified accuracy of convergence guarantees the unitarity of T-matrix [9].

Equation (21) describes scattering on weekly bound system $\left(\mathrm{Re}\left(z\right)\gg E_{\min}^{B}\right)$. According to the approximation of Eq. (15), equation (21) corresponds to the processes in which single collision dominate. The expression of the kernel of the obtained asymptotic equations with the Green's function G_1 indicates that these equations are adequate for elastic and quasi-elastic scattering reactions (the internal energy of the system does not change)

Let us insert Eq. (23) in Eq. (6):

$$T\left(z\right)=\sum_{\alpha}T_{\alpha}\left(z\right)\left(1+G_{1}\left(z\right)\sum_{\beta\neq\alpha}T_{\beta}\left(z\right)\right),\quad\mathrm{Re}\left(z\right)\gg E_{\min}^{B}, \tag{24}$$

$$\alpha,\beta\equiv\left(mn\right),\quad m,n=1,2,\ \ldots,N;\ \ m<n,N\geq3.$$

Equation (24) is a particular unitary solution of Eq. (16), which at the same time represents an asymptotic solution of Eqs. (2) and (9).

Thus, by using the Heitler formalism in multi-particle problem we got multi-particle asymptotic equations. In particular, keeping in the right-hand side of Eq. (9) the high-energy terms we obtained asymptotic equation (16). Based on this equation we constructed the Faddeev type compact the system of N-particle integral equations in asymptotic, Eq. (21). Using the convergence of the iterative series corresponding to Eq. (21) we obtained Eq. (24), the unitary asymptotic solution of multi-particle Eqs. (2) and (9), which is identical to the result proposed by us earlier [9].

Prof. Alexander Kvinikhidze and Jambul Gegelia provided systematic assistance during the work on the article. We express our gratitude to Prof. Anzor Khelashvili and Ilia Lomidze. Also we thenk Mikheil Makhviladze for his support.

Appendix

Below we establish the high-energy behavior of operators determined by Eqs. (8) and (13). The obtained result is used for the estimation of the kernel of multi-particle equations.

Let as show that

$$T_\alpha(z) \approx \upsilon_\alpha, \quad \mathrm{Re}(z) > \mathrm{Re}(\bar{z}). \tag{25}$$

Here $z = E_0 + i\varepsilon$. E_0 is a kinetic energy of the considered system. $\bar{z} = \bar{E} + i\varepsilon$. $\bar{E}$ is the fixed energy, which satisfies the condition: $\bar{E} \gg E^B_{\min}$. $E^B_{\min}$ denotes the absolute value of the minimum of all allowed binding energies in the considered system.

Using the symbolic notation $\alpha \equiv (mn)$, we write the connection among t_α and T_α operators:

$$\left\langle q'_1\ q'_2 . .\ q'_N \left| T_\alpha(z) \right| q_1 q_2 . .\ q_N \right\rangle = \prod_{l \neq m \neq n \neq l}^{N} \delta\left(q'_l - q_l\right) \left\langle q'_m\ q'_n \left| t_\alpha\left(z'\right) \right| q_m q_n \right\rangle, \tag{26}$$

where the pure two particle interaction operator t_α satisfies the LS integral equation:

$$t_\alpha\left(z'\right) = \upsilon_\alpha + \upsilon_\alpha g_0\left(z'\right) t_\alpha\left(z'\right). \tag{27}$$

Here g_0 is the Greenq's function of two free particles.

$$z' = E' + i\varepsilon, \quad E' = E_0 - \sum_{n \neq l \neq m} \frac{q_l^2}{2m_l}. \tag{28}$$

E_0 and q_l are fixed parameters. According to the Klein- Zemach theorem [10] we have:

$$t_\alpha\left(z'\right) \approx \upsilon_\alpha, \quad \mathrm{Re}\left(z'\right) > \mathrm{Re}(\bar{z}). \tag{29}$$

Inserting the relation (29) into Eq. (26) we get Eq. (25).

Let us express T_α from Eq. (8) and substitute the obtained result in Eq. (25):

$$T_\alpha(z) = \left(1 - \upsilon_\alpha G_0(z)\right)^{-1} \upsilon_\alpha \approx \upsilon_\alpha, \quad \mathrm{Re}(z) > \mathrm{Re}(\bar{z}). \tag{30}$$

Due to Eqs. (25) and (30) we obtained:

$$\left\| T_\alpha(z) G_0(z) \right\| \approx \left\| \upsilon_\alpha G_0(z) \right\| \approx \varepsilon, \quad \mathrm{Re}(z) > \mathrm{Re}(\bar{z}). \tag{31}$$

Similarly to Eq. (26) we write:

$$\left\langle q'_1\ q'_2 . .\ q'_N \left| K_\alpha(z) \right| q_1 q_2 . .\ q_N \right\rangle = \prod_{l \neq m \neq n \neq l}^{N} \delta\left(q'_l - q_l\right) \left\langle q'_m\ q'_n \left| k_\alpha\left(z'\right) \right| q_m q_n \right\rangle. \tag{32}$$

Here k_α is a two particle Hermitian operator:

$$k_\alpha\left(z'\right) = \upsilon_\alpha + \upsilon_\alpha g_2\left(z'\right) k_\alpha\left(z'\right), \tag{33}$$

where g_2 can be expressed as follows:

$$g_2\left(z'\right) = \frac{1}{2}\left(g_0\left(z'\right) + g_0\left(\tilde{z}'\right)\right). \tag{34}$$

The parameter z' in Eq. (34) is defined from equalities (28).

Due to the mentioned Klein-Zemach theorem, one can show similarly to Eq. (25):

$$K_\alpha(z) \approx \upsilon_\alpha, \quad \mathrm{Re}(z) > \mathrm{Re}(\bar{z}). \tag{35}$$

According to Eqs. (13) and (35) we have:

$$\left\| K_\alpha(z) G_2(z) \right\| \approx \left\| \upsilon_\alpha G_2(z) \right\| \approx \varepsilon, \quad \mathrm{Re}(z) > \mathrm{Re}(\bar{z}). \tag{36}$$

From Eq. (19) in the approximation of Eq. (35) we obtain [8]:

$$T_\alpha(z) = \upsilon_\alpha + \upsilon_\alpha G_1(z) T_\alpha(z). \tag{37}$$

Using Eqs. (37) and (25) one can write:

$$\left\| T_\alpha(z) G_1(z) \right\| \approx \left\| \upsilon_\alpha G_1(z) \right\| \approx \varepsilon, \quad \mathrm{Re}(z) > \mathrm{Re}(\bar{z}). \tag{38}$$

Thus, it is shown that in the asymptotic $\left(\mathrm{Re}(z) \gg E^B_{\min}\right)$, the kernels of multi-particle equations are sufficiently small, so that the corresponding iterative series converge.

References

[1] L. D. Faddeev, Zh. Exp. Teor. Fiz.. 1960, 39, 1945.

[2] O. A. Yakubovsky, Sov. J. Nucl. Phys. 1967, Vol. 5, p. 1312
[Yad. Fiz. 5, 1312 (1967)].

[3] E. O. Alt, P. Grassberger, W. Sandhas. Nucl. Physics. 1967, Vol. B2, p.167,
A. A. Khelashvili, Preprint JINR, P2-3371,Dubna, 1967.

[4] S. P. Merkuryev, L. D. Faddeev. Quantum scattering theory for the system of a few particles (in Russian. Moscow, "Nauka", 1985).

[5] J. V. Mebonia and T. I. Kvarackheliya, Physics Letters. 1980, Vol. 90B, num. 1,2, p. 17.

[6] Mavrin L. Goldberger, Kenneth M. Watson. Collision theory. Jonh Wiley and Sons, Inc. New York-London-Sydney 1964.

[7] J. V. Mebonia, Physics Letters. 1969, Vol. 30B, num. 3, p. 153.

[8] http://www.phy.ohiou.edu/∼elster/qm1_12.pdf.

[9] Vagner Jikia, Jemal Mebonia. On Approximate Asymptotic Solution of Integral Equations of the Collision Theory. ArXiv: 1312. 0865vl [math-ph] 3 Dec 2013.

[10] Michael Reed, Berry Simon. Methods of modern mathematical physics. Vol. 3. Scattering theory. New York, San Francisco, London. Academic press. 1979,
A. Klein, Ch. Zemach. Analytic properties of the amplitude for the scattering of a particle by a central potential. Vol. 7, Issue 4, pp. 440-455, August 1959.